

www.tajpharma.com

Tamoxifen as breast cancer treatment.

Adylaro® Bleomycin is an anti-cancer ("antineoplastic" or "cytotoxic") chemotherapy drug. Bleomycin is classified as an "antitumor antibiotic." Fever of 100.4° F (38° C) or higher, chills (possible signs of infection).

Adylaro
(BLEOMYCIN)

Adylaro® Bleomycin for Injection USP 15 UNITS

TREATMENT OF SQUAMOUS CELL CANCERS, MELANOMA, SARCOMA, TESTICULAR AND OVARIAN CANCER, HODGKIN'S, AND NON-HODGKIN'S LYMPHOMA.

How Bleomycin Is Given:

As an infusion into the vein (intravenous, IV).

As an intrapleural - into the pleural cavity (the space between the lining of the lung and the chest wall) injection. This is done to treat a malignant pleural effusion (buildup of fluid). Intrapleural chemotherapy is used primarily for symptom relief.

THERE IS NO PILL FORM OF BLEOMYCIN.

YOU MAY RECEIVE A TEST DONE OF BLEOMYCIN BASED ON YOUR CONDITION.

Adylaro
(BLEOMYCIN)

What Bleomycin Is Used For: Used in the treatment of squamous cell cancers, melanoma, sarcoma, testicular and ovarian cancer, Hodgkin's, and non-Hodgkin's lymphoma. Also used to treat malignant pleural effusion (buildup of fluid in the space between the lining of the lung and the chest wall).

What Bleomycin Is Used For:

Used in the **treatment of squamous cell cancers, melanoma, sarcoma, testicular and ovarian cancer, Hodgkin's, and non-Hodgkin's lymphoma.**

Also used to treat malignant pleural effusion (buildup of fluid in the space between the lining of the lung and the chest wall).

Note: If a drug has been approved for one use, physicians may elect to use this same drug for other problems if they believe it may be helpful.

DO NOT THROW AWAY ANY MEDICINES VIA WASTEWATER OR HOUSEHOLD WASTE. ASK YOUR PHARMACIST HOW TO THROW AWAY MEDICINES YOU NO LONGER USE. THESE MEASURES WILL HELP PROTECT THE ENVIRONMENT.

How Bleomycin Is Given:

As an infusion into the vein (intravenous, IV).

As an intrapleural - into the pleural cavity (the space between the lining of the lung and the chest wall) injection. This is done to treat a malignant pleural effusion (buildup of fluid).

Intrapleural chemotherapy is used primarily for symptom relief.

There is no pill form of bleomycin.

You may receive a test done of bleomycin based on your condition.

The amount of bleomycin that you will receive and the method it is given depends on many factors, including your height and weight, your general health or other health problems, and the type of cancer or condition being treated. Your doctor will determine your dose, schedule, and how it will be given.

Monitoring and Testing While Taking Bleomycin:

You will be checked regularly by your health care professional while you are taking bleomycin, to monitor side effects and check your response to therapy. Periodic blood work to monitor your complete blood count (CBC) as well as the function of other organs (such as your kidneys and liver) will also be ordered by your doctor.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma™ India Product

TAJ
PHARMA
A TAJ Enterprise

Your doctor will determine your dose, schedule, and how it will be given.