

www.tajpharma.com

Take CAPRIZANT® just once a day for symptom relief you can count on. The recommended dose of CAPRIZANT for the treatment of moderate-to-severe Restless Legs Syndrome (RLS) in adults is 600 mg taken once a day with food in the evenings (usually at around 5 PM). Do not cut, crush, or chew your tablets.

  
**Caprizant®**  
Gabapentin enacarbil  
Extended-Release Tablet  
300 MG & 600 MG

Do not drink alcohol while taking CAPRIZANT because it may increase the risk of side effects.  
Treatment for: Restless Legs Syndrome, Postherpetic Neuralgia  
CAPRIZANT® (gabapentin enacarbil) is a gabapentin prodrug indicated for the once-daily treatment of restless legs syndrome and postherpetic neuralgia.



**Caprizant®**  
Gabapentin enacarbil  
Extended-Release Tablet  
300 MG & 600 MG

**CAPRIZANT® (gabapentin enacarbil) is a prescription medicine used to treat adults with moderate-to-severe primary RLS. When used for RLS, the most common side effects of CAPRIZANT include sleepiness and dizziness.**

CAPRIZANT® (gabapentin enacarbil) Extended-Release Tablets

#### INDICATIONS

CAPRIZANT® (gabapentin enacarbil) is a prescription medicine used to: treat adults with moderate to severe primary Restless Legs Syndrome (RLS). CAPRIZANT is not for people who need to sleep during the daytime and stay awake at night. manage pain from damaged nerves (postherpetic neuralgia) that follows healing of shingles (a painful rash that comes after a herpes zoster infection) in adults.

#### IMPORTANT SAFETY INFORMATION

Do not drive after taking your dose of CAPRIZANT until you know how it affects you, including the morning after you take it. Do not operate heavy machinery or do other dangerous activities until you know how CAPRIZANT affects you. CAPRIZANT can cause sleepiness, dizziness, slow thinking, and can affect your coordination. Ask your healthcare provider when it is okay to do these activities. Do not take other medicines that make you sleepy or dizzy while taking CAPRIZANT without talking to your healthcare provider. Taking CAPRIZANT with these other medicines may make your sleepiness or dizziness worse. With CAPRIZANT, you can get control of your RLS symptoms and get the rest you need.

CAPRIZANT can offer relief when RLS symptoms affect you most—during the night.

Is your moderate-to-severe primary Restless Legs Syndrome (RLS) leaving you on edge? It's hard enough living with the uncomfortable feelings in your legs, but the urge to get up and move can make it nearly impossible to rest. When used for RLS, the most common side effects of CAPRIZANT include sleepiness and dizziness. CAPRIZANT (gabapentin enacarbil) Extended-Release Tablets for oral use 300mg or 600mg Tablets.

**Food and Drug Administration (FDA) approved .Prescription Only (POM)**

A Taj Pharma™ India Product