

www.tajpharma.com

This medication is used to treat certain types of cancer (such as leukemia, lymphoma).

Tresicta® Chlorambucil oral tablet is a prescription medication used to treat certain types of cancer of the blood and lymph nodes.

TRESICTA®
chlorambucil | 2mg/ 5mg
tablets



TRESICTA®
chlorambucil | 2mg/ 5mg
tablets



STARTING DOSES OF 6-12 MG DAILY UNTIL LEUCOPENIA OCCURS ARE RECOMMENDED FOLLOWED BY 2-8 MG DAILY INDEFINITELY.
Store between 2°C and 8°C in a dry place.
WARNING : Tresicta®(chlorambucil) can severely suppress bone marrow function. Chlorambucil is a carcinogen in humans. Chlorambucil is probably mutagenic and teratogenic in humans. Chlorambucil produces human infertility

Tresicta® Chlorambucil contains a medicine called chlorambucil. This belongs to a group of medicines called cytotoxics (also called chemotherapy). Chlorambucil is used to **treat some types of cancer and certain blood problems.** It works by reducing the number of abnormal cells your body makes.

Chlorambucil is used for:

- Hodgkin's disease and Non-Hodgkin's Lymphoma. Together, these form a group of diseases called lymphomas. They are cancers formed from cells of the lymphatic system.
 - Chronic lymphocytic leukaemia. A type of white blood cell cancer where the bone marrow produces a large number of abnormal white cells.
 - Waldenstrom's macroglobulinaemia A rare lymphoma associated with an uncontrolled increase of B-cells, a type of white blood cell, resulting in the release of an abnormal protein into the blood.
- Ask your doctor if you would like more explanation about these diseases.

You must talk to a doctor if you do not feel better or if you feel worse.

Do not take Chlorambucil :

If you are allergic (hypersensitive) to chlorambucil or any of the other ingredients of Chlorambucil tablets (See section 6).
If you are not sure, talk to your doctor or nurse before taking Chlorambucil.

Warnings and precautions

Before you take Chlorambucil, tell your doctor or nurse if:

you have been recently vaccinated, or planning to be vaccinated with a live vaccine (see Other medicines and Chlorambucil). Chlorambucil can make your body less able to fight infections.
you are a potential candidate for bone marrow transplantation (autologous stem cell transplantation) as the long term use of Chlorambucil may reduce the amount of stem cells.

you have had radiotherapy or chemotherapy, now or recently
you have a liver or kidney problems
you have a kidney problem (nephrotic syndrome), had high pulse dosing regimen or ever had a fit or convulsion.
If you had ever had fits or convulsions, you might be more at risk of having fits or convulsions when taking Chlorambucil.
It is possible that the use of Chlorambucil, particularly long term use, may increase the risk of developing a secondary blood cancer. In many cases, patients who develop this have also received another type of chemotherapy or some form of radiation therapy. Symptoms of a secondary blood cancer include tiredness, fever, infection and bruising.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma™ India Product

DO NOT THROW AWAY ANY MEDICINES VIA WASTEWATER OR HOUSEHOLD WASTE. ASK YOUR PHARMACIST HOW TO THROW AWAY MEDICINES YOU NO LONGER USE. THESE MEASURES WILL HELP PROTECT THE ENVIRONMENT.

TAJ
PHARMA®
A TAJ Enterprise

Chlorambucil should be taken on an empty stomach.