

**Desloratadine 5mg film-coated tablets Taj Pharma**

**Package leaflet: Information for the patient**

**Desloratadine 5mg film-coated tablets**

**desloratadine**

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Desloratadine is and what it is used for
2. What you need to know before you take Desloratadine
3. How to take Desloratadine
4. Possible side effects
5. How to store Desloratadine
6. Contents of the pack and other information

**1. What Desloratadine is and what it is used for**

Desloratadine is an antiallergy medicine that does not make you drowsy. It helps control your allergic reaction and its

symptoms.

Desloratadine is indicated for adults and adolescents (12 years of age and older) to: relieve symptoms associated with allergic rhinitis (inflammation of the nasal passages caused by an allergy, for example, hay fever or allergy to dust mites). These symptoms include sneezing, runny or itchy nose, itchy palate, and itchy, red or watery eyes.

Desloratadine is also used to relieve the symptoms associated with urticaria (a skin condition caused by an allergy). These symptoms include itching and hives.

Relief of these symptoms lasts a full day and helps you to resume your normal daily activities and sleep.

You must talk to a doctor if you do not feel better or if you feel worse.

**What you need to know before you take Desloratadine Do not take Desloratadine:**

- if you are allergic to desloratadine, to any of the other ingredients of this medicine (listed in section 6) or to loratadine.

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Desloratadine if you:

- have medical or familial history of seizures.
- have poor kidney function.

If this applies to you, or if you are not sure, please check with your doctor or pharmacist.

**Children**

Desloratadine tablets are not suitable for children under 12 years of age.

#### **Other medicines and Desloratadine**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

#### **Desloratadine with alcohol**

Caution is advised when taking desloratadine with alcohol.

#### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

If you are pregnant or breast-feeding a baby, taking Desloratadine is not recommended.

#### **Driving and using machines**

At the recommended dose, Desloratadine is not expected to cause you to be drowsy or less alert. However, very rarely some people experience drowsiness, which may affect their ability to drive or use machines.

#### **Desloratadine contains Sunset**

**Yellow Aluminium Lake** which may cause allergic reactions.

## **2. How to take Desloratadine**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose for adults and adolescents (12 years of age and older) is one tablet once a day. Swallow the tablet

whole with water, with or without food.

Regarding the duration of treatment, your doctor will determine the type of allergic rhinitis you are suffering from and will determine for how long you should take Desloratadine.

If your allergic rhinitis is intermittent (presence of symptoms for less than 4 days per week or for less than 4 weeks), your doctor will recommend you a treatment schedule that will depend on the evaluation of the history of your disease.

If your allergic rhinitis is persistent (presence of symptoms for 4 days or more per week and for more than 4 weeks), your doctor may recommend you a longer term treatment.

For urticaria, the duration of treatment may be variable from patient to patient and therefore you should follow the instructions of your doctor.

#### **If you take more Desloratadine than you should**

Take Desloratadine only as it is prescribed for you. No serious problems are expected with accidental overdose. However, if you take more Desloratadine than you were told to, contact your doctor or pharmacist.

#### **If you forget to take Desloratadine**

If you forget to take your dose on time, take it as soon as possible, then go back to your regular dosing schedule. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **3. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. In adults, side effects were about the same as with a dummy tablet. However, fatigue, dry mouth and headache were reported more often than with a dummy tablet. In adolescents, headache was the most commonly reported side effect.

**If you notice any of the following side effects, stop taking this medicine and contact your doctor or go to the nearest hospital casualty department straight away:**

**Very rare: (may affect up to 1 in 10,000 people):**

- Severe allergic reactions such as difficulty in breathing, shortness of breath, wheezing, itching, hives and swelling of the face, lips, tongue or other parts of the body and rash
- Fits (seizures)
- Liver disease (nausea, vomiting, loss of appetite, feeling generally unwell, fever, itching, yellowing of the skin and eyes, light coloured bowel motions, dark coloured urine)

**Not known: (frequency cannot be estimated from available data):**

- A change in the way the heart beats, which may make you feel dizzy or faint. This may be seen in tests of the electrical activity of the heart ('electrocardiogram' or ECG)

In clinical studies with desloratadine, the following side effects were reported as:

**Common: (may affect up to 1 in 10 people):**

- Fatigue
- Dry mouth
- Headache

During the marketing of desloratadine, the following side effects were reported in

adults, as:

**Very rare: (may affect up to 1 in 10,000 people):**

- Fast heartbeat
- Being sick (vomiting)
- Dizziness
- Muscle pain
- Restlessness with increased body movement
- Stomach ache
- Upset stomach
- Drowsiness
- Seeing, feeling or hearing things that are not there (Hallucinations)
- Pounding or irregular heartbeat
- Feeling sick (nausea)
- Diarrhoea
- Inability to sleep
- Abnormal liver function tests

**Not known: (frequency cannot be estimated from available data):**

- Abnormal behavior
- Aggression
- Unusual weakness
- Increased sensitivity of the skin to the sun, even in the case of hazy sun, and to UV light, for instance to UV lights of a solarium
- Weight increased, increased appetite

**Additional side effects in children and adolescents**

**Not known: (frequency cannot be estimated from available data):**

- Slow heart beat

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### 4. How to store Desloratadine

Keep this medicine out of the sight and reach of children. This medicine does not require any special storage conditions.

Do not use this medicine after the expiry date, which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### 5. Contents of the pack and other information

##### What Desloratadine contains

The active substance is desloratadine. Each tablet contains 5mg of desloratadine.

The other ingredients of the tablet are: magnesium stearate, sodium lauril sulfate, silica colloidal anhydrous, microcrystalline cellulose and pregelatinised maize starch.

The tablet film coating contains: poly (vinyl alcohol), indigo carmine aluminium lake, sunset yellow aluminium lake (see section 2 Desloratadine contains sunset yellow), macrogol, talc and titanium dioxide.

#### 6. What Desloratadine looks like and contents of the pack

Desloratadine 5mg film-coated tablets are packed in blisters in packs of 2, 3, 5, 7, 10, 15,

20, 30, 50, 60, 90 and 100 tablets.

Not all pack sizes may be marketed.

#### 7. Manufactured By: Taj Pharmaceuticals Ltd.

at: Plot. No. 220, Mahagujarat  
Industrial Estate, At & Post-Moraiya,  
Tal-Sanand, Dist- Ahmedabad Gujarat (India)