

## **Esomeprazole Sodium for Injection 40mg Taj Pharma**

### Package Leaflet: Information for the patient

Esomeprazole 40 mg powder for solution for injection/infusion

**Read all of these leaflet carefully before you are given this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet?**

1. What Esomeprazole is and what it is used for
2. What you need to know before you are given Esomeprazole
3. How Esomeprazole is given
4. Possible side effects
5. How to store Esomeprazole
6. Contents of the pack and other information

#### **1. What Esomeprazole is and what it is used for**

Esomeprazole contains a medicine called esomeprazole. This belongs to a group of medicines called ‘proton pump inhibitors’. They work by reducing the amount of acid that your stomach produces.

Esomeprazole is used for the short-term treatment of certain conditions, when you are unable to have treatment by mouth. It is used to treat the following conditions:

#### Adults

- ‘Gastro-oesophageal reflux disease’ (GERD). This is where acid from the stomach escapes into the gullet (the tube which connects your throat to your stomach) causing pain, inflammation and heartburn.
- Stomach ulcers caused by medicines called NSAIDs (Non-Steroidal Anti-Inflammatory Drugs). Esomeprazole can also be used to stop stomach ulcers from forming if you are taking NSAIDs.
- Prevention of rebleeding in adults following therapeutic endoscopy for acute bleeding gastric or duodenal ulcers.

#### Children and adolescents aged 1-18 years

- ‘Gastro-oesophageal reflux disease’ (GERD). This is where acid from the stomach escapes into the gullet (the tube which connects your throat to your stomach) causing pain, inflammation and heartburn.

#### **2. What you need to know before you are given Esomeprazole**

##### **You must not be given Esomeprazole if**

- you are allergic to esomeprazole or any of the other ingredients of this medicine (listed in section 6)

- you are allergic to other proton pump inhibitor medicines (e.g. pantoprazole, lansoprazole, rabeprazole, omeprazole)
- you are taking a medicine containing nelfinavir (used to treat HIV infection).

You must not be given Esomeprazole if any of the above apply to you. If you are not sure, talk to your doctor before you are given this medicine.

### Warnings and Precautions

Talk to your doctor, pharmacist or nurse before you are given Esomeprazole if:

- you have severe liver problems
- you have severe kidney problems.
- you have ever had a skin reaction after treatment with a medicine similar to Esomeprazole that reduces stomach acid.
- you are due to have a specific blood test (Chromogranin A).

Esomeprazole may hide the symptoms of other diseases. **Therefore, if any of the following happen to you before you are given Esomeprazole or after you are given it, talk to your doctor straight away:**

- you lose a lot of weight for no reason and have problems swallowing
- you get stomach pain or indigestion
- you begin to vomit food or blood

- you pass black stools (blood-stained faeces).

Taking a proton pump inhibitor like Esomeprazole, especially over a period of more than one year, may slightly increase your risk of fracture in the hip, wrist or spine. Tell your doctor if you have osteoporosis or if you are taking corticosteroids (which can increase the risk of osteoporosis).

If you get a rash on your skin, especially in areas exposed to the sun tell your doctor as soon as you can, as you may need to stop your treatment with Esomeprazole. Remember to also mention any other ill-effects like pain in your joints.

### Other medicines and Esomeprazole

Tell your doctor if you are taking, have recently taken or might take any other medicines. This includes medicines that you buy without a prescription. This is because Esomeprazole can affect the way some medicines work and some medicines can have an effect on Esomeprazole.

You must not be given Esomeprazole if you are taking a medicine containing **nelfinavir** (used to treat HIV infection).

Tell your doctor or nurse if you are taking any of the following medicines:

- atazanavir (used to treat HIV infection)
- clopidogrel (used to prevent blood clots)
- ketoconazole, itraconazole or voriconazole (used to treat infections caused by a fungus)
- erlotinib (used to treat cancer)
- citalopram, imipramine or clomipramine (used to treat depression)

- diazepam (used to treat anxiety, relax muscles or in epilepsy)
- phenytoin (used in epilepsy). If you are taking phenytoin, your doctor will need to monitor you when you start or stop having Esomeprazole.
- medicines that are used to thin your blood, such as warfarin. Your doctor may need to monitor you when you start or stop having Esomeprazole.
- Cilostazol (used to treat intermittent claudication – a pain in your legs when you walk which is caused by an insufficient blood supply)
- cisapride (used for indigestion and heartburn)
- digoxin (used for heart problems)
- methotrexate (a chemotherapy medicine used in high doses to treat cancer) – if you are taking a high dose of methotrexate, your doctor may temporarily stop your Esomeprazole treatment.
- tacrolimus (organ transplantation)
- rifampicin (used for treatment of tuberculosis)
- St. John's wort (*Hypericum perforatum*) (used to treat depression).

### **Pregnancy and breast-feeding**

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before you are given this medicine. Your doctor will decide whether you can be given Esomeprazole during this time.

It is not known if Esomeprazole passes into breast milk. Therefore, you should not be given Esomeprazole if you are breast-feeding.

### **Driving and using machines**

Esomeprazole is not likely to affect you being able to drive or use any tools or machines. However, side effects such as dizziness and blurred vision may uncommonly occur (see section 4). If affected, you should not drive or use machines.

### **Esomeprazole contains sodium**

This medical product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially “sodium-free”.

### **3. How Esomeprazole is given**

Esomeprazole can be given to children and adolescents aged 1-18 years and adults including the elderly.

### **Being given Esomeprazole**

#### **Adults**

- Esomeprazole will be given to you by your doctor who will decide how much you need.
- the recommended dose is 20 mg or 40 mg once a day
- if you have severe liver problems, the maximum dose is 20 mg a day (GERD)
- the medicine will be given to you as an injection or infusion into one of your veins. This will last for up to 30 minutes
- the recommended dose for prevention of re-bleeding gastric or duodenal ulcers, is 80 mg administered as intravenous infusion over 30 minutes followed by a continuous infusion of 8 mg/hr given over 3 days. If you have severe liver problems, a continuous infusion of 4 mg/hr given over 3 days may be sufficient.

### **Use in children and adolescents aged 1-18 years**

- Esomeprazole will be given by your doctor who will decide how much you need.
- For children 1-11 years, the recommended dose is 10 or 20 mg given once a day.
- For children 12-18 years, the recommended dose is 20 or 40 mg given once a day.
- The medicine will be given as an injection or infusion into a vein. This will last up to 30 minutes.

### **If you are given too much Esomeprazole**

If you think you have been given too much Esomeprazole, talk to your doctor straight away.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

### **If you notice any of the following serious side effects, stop taking Esomeprazole and contact a doctor immediately:**

- sudden wheezing, swelling of your lips, tongue and throat or body, rash, fainting or difficulties in swallowing (severe allergic reaction)
- reddening of the skin with blisters or peeling. There may also be severe blisters and bleeding in the lips, eyes, mouth, nose and genitals. This could be 'Stevens-Johnson syndrome' or 'toxic epidermal necrolysis'.

- yellow skin, dark urine and tiredness which can be symptoms of liver problems.

These effects are rare and may affect up to 1 in 1,000 people.

Esomeprazole may in very rare cases affect the white blood cells leading to immune deficiency. If you have an infection with symptoms such as fever with a **severely** reduced general condition or fever with symptoms of a local infection such as pain in the neck, throat or mouth or difficulties in urinating, you must consult your doctor as soon as possible so that a lack of white blood cells (agranulocytosis) can be ruled out by a blood test. It is important for you to give information about your medication at this time.

Other side effects include:

### **Common (may affect up to 1 in 10 people)**

- headache
- effects on your stomach or gut: diarrhoea, stomach pain, constipation, wind (flatulence)
- feeling sick (nausea) or being sick (vomiting)
- injection site reaction
- benign polyps in the stomach.

### **Uncommon (may affect up to 1 in 100 people)**

- swelling of the feet and ankles
- disturbed sleep (insomnia)
- dizziness, tingling feelings such as "pins and needles", feeling sleepy
- spinning feeling (vertigo)

- eyesight problems such as blurred vision
- dry mouth
- changes in blood tests that check how the liver is working
- skin rash, lumpy rash (hives) and itchy skin
- fracture of the hip, wrist or spine (if Esomeprazole is used in higher doses and over long duration).

**Rare (may affect up to 1 in 1,000 people)**

- blood problems such as a reduced number of white cells or platelets. This can cause weakness, bruising or make infections more likely.
- low levels of sodium in the blood. This may cause weakness, being sick (vomiting) and cramps.
- feeling agitated, confused or depressed
- taste changes
- suddenly feeling wheezy or short of breath (bronchospasm)
- an inflammation of the inside of the mouth
- an infection called “thrush” which can affect the gut and is caused by a fungus
- liver problems, including jaundice which can cause yellow skin, dark urine, and tiredness
- hair loss (alopecia)
- skin rash on exposure to sunshine
- joint pains (arthralgia) or muscle pains (myalgia)
- generally feeling unwell and lacking energy
- increased sweating.

**Very rare (may affect up to 1 in 10,000 people)**

- changes in blood count including agranulocytosis (lack of white blood cells)
- aggression
- seeing, feeling or hearing things that are not there (hallucinations)
- severe liver problems leading to liver failure and inflammation of the brain
- sudden onset of a severe rash or blistering or peeling skin. This may be associated with a high fever and joint pains (Erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis).
- muscle weakness
- severe kidney problems
- enlarged breasts in men.

**Not known (frequency cannot be estimated from the available data)**

- If you are on Esomeprazole for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary muscle contractions, disorientation, convulsions, dizziness, increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium.
- Inflammation in the gut (leading to diarrhoea).
- Rash, possibly with pain in the joints.



### **Reporting of side effects**

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet.

### **5. How to store Esomeprazole**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton or vial after EXP. The expiry date refers to the last day of that month.

Store below 30°C. Store in the original package in order to protect from light.

Chemical and physical in-use stability data has been demonstrated for 12 hours at 30°C. From microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2-8°C, unless reconstitution has taken place in controlled and aseptic conditions.

Do not use this medicine if you notice that the visual appearance has changed (only clear solution should be used).

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

### **6. Contents of the pack and other information**

#### **What Esomeprazole contains**

- The active substance is esomeprazole sodium. Each vial contains 42.6 mg of esomeprazole sodium, equivalent to 40 mg of esomeprazole.
- The other ingredients are disodium edetate and sodium hydroxide (for pH adjustment).

#### **What Esomeprazole looks like and contents of the pack**

Esomeprazole is a white to off-white powder. It comes in a 5 ml colorless glass vial with grey rubber stopper and grey flip off aluminum seal containing 40 mg powder for solution for injection/infusion.

Esomeprazole is available in the following pack sizes:

- pack with 1 vial
- pack with 1x10 vials.

Not all pack sizes may be marketed.

#### **7. Manufactured in India by: TAJ PHARMACEUTICALS LTD.**

Mumbai, India

Unit No. 214.Old Bake House,  
Maharashtra chambers of Commerce Lane,

Fort, Mumbai - 400001

at:Gujarat, INDIA.

Customer Service and Product Inquiries:

1-800-TRY-FIRST (1-800-222-434 & 1-800-222-825)

Monday through Saturday 9:00 a.m. to 7:00 p.m. EST

E-mail: tajgroup@tajpharma.com