

Imipramine HCl Tablets 25mg USP Taj Pharma

(Imipramine)

Patient Leaflet: Information for the user

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What Imipramine tablets are and what they are used for
2. Before you take Imipramine
3. How to take Imipramine
4. Possible side effects
5. How to store Imipramine
6. Further information

1. What Imipramine tablets are and what they are used for

Imipramine belongs to a group of medicines called tricyclic antidepressant drugs. These medicines alter the levels of chemicals in the brain to relieve the symptoms of depression. Imipramine is used:

- to treat the symptoms of depression.
- for the relief of bed-wetting at night by children.

2. Before you take Imipramine

Do not take Imipramine tablets and **tell** your doctor if you or your child (if they are the patient):

- are **allergic** (hypersensitive) to imipramine, other tricyclic antidepressants or any of the other ingredients (see section 6).
- have **heart disease** such as irregular heart beats, heart block or have recently had a heart attack
- suffer from periods of increased and exaggerated behaviour (**mania**)
- have **severe liver** disease
- suffer with **porphyria** (a genetic disorder of the red blood cells haemoglobin causing skin blisters, abdominal pain and brain/nervous system disorders)
- are **not able to pass water**
- have increased pressure in the eye (**glaucoma**)
- are taking monoamine oxidase inhibitors (**MAOI**) or you have taken MAOIs within the previous 14 days for depression
- if the **child is under 6** years old.

Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- If you have previously had thoughts about killing or harming yourself.
- If you are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in young adults (less than 25 years old) with psychiatric conditions who were treated with an antidepressant.

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away.**

You may find it helpful to tell a relative or close friend that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

Check with your doctor or pharmacist before taking

Imipramine tablets if you or your child (if they are the patient):

- have any **psychiatric disorder** (eg schizophrenia or manic depression)
- are **withdrawing** from **alcohol or medicines** used to treat fits
- have ever **had glaucoma** or an **enlarged prostate gland**
- have an **overactive thyroid** gland and are taking medicines to treat a thyroid disorder
- have a history of **epilepsy** or **brain damage**
- have **low blood pressure** or **poor circulation**
- have **severe kidney** disease
- have a **tumour** of the adrenal gland (eg pheochromocytoma or neuroblastoma)
- suffer from **panic attacks**
- suffer from long term **constipation**
- wear **contact lenses**
- are being given electroconvulsive therapy (**ECT**)
- are due to have **any surgery**, including dental, that **involves anaesthetic**.

Taking other medicines

Please **tell your doctor or pharmacist** if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Especially:

- medicines to treat epilepsy such as barbiturates, phenytoin, carbamazepine, phenobarbital
- medicines called “benzodiazepines” such as diazepam, nitrazepam, oxazepam, alprazolam
- medicines to treat depression, such as selective serotonin reuptake inhibitors (SSRIs) such as fluoxetine, fluvoxamine
- disulfiram to treat alcohol addiction
- nicotine replacement therapy
- methylphenidate (used to treat attention deficit/hyperactivity disorder (ADHD))
- medicines to stop your blood clotting (eg warfarin)
- antihistamines (medicines to treat allergies)
- altretamine (to treat some types of cancer)
- apraclonidine and brimonidine (to treat glaucoma)
- baclofen (a muscle relaxant)
- painkillers such as nefopam, tramadol, codeine, dihydrocodeine
- medicines to treat some heart conditions such as diltiazem, verapamil, labetalol, propranolol, quinidine
- medicines to treat angina that you spray or dissolve under your tongue (eg glyceryl trinitrate “GTN”, isosorbidedinitrate)
- any medicines to treat high blood pressure such as guanethidine, debrisoquine, bethanidine, methyldopa, reserpine, clonidine or diuretics (“water” tablets)
- medicines to treat some mental illnesses such as thioridazine, chlorpromazine
- entacapone or selegiline (to treat Parkinson’s disease)
- oral contraceptives (“the pill”) or hormone replacement therapy (HRT)
- appetite suppressants
- sympathomimetic medicines such as adrenaline (epinephrine), ephedrine, isoprenaline, noradrenaline (norepinephrine), phenylephrine and phenylpropanolamine (these may be present in many cough and cold remedies or local anaesthetics)

- ritonavir (to treat HIV).

Pregnancy and breast-feeding

Imipramine tablets should not be taken during pregnancy or if breast-feeding. If Imipramine tablets are taken in the last 3 months the baby may be born with breathing difficulties, lethargy, colic, irritability, changes in blood pressure, tremors, spasm. Imipramine tablets should be withdrawn at least 7 weeks before the expected delivery date.

Driving and using machines

Imipramine may impair your alertness or cause drowsiness or blurred vision, alcohol can make these symptoms worse. Make sure you are not affected before you drive or operate machinery.

Blood tests

Whilst taking Imipramine tablets your doctor will regularly monitor your blood cell levels or liver function.

Dental check ups

As Imipramine tablets can cause problems with your teeth, it is advisable to have regular dental checks ups.

Sugar intolerance

If you have been told you have an intolerance to some sugars, contact your doctor before taking this medicine, as it contains types of sugars called lactose or sucrose.

3. How to take Imipramine

Always take Imipramine tablets exactly as your doctor has told you. If you are not sure, check with your doctor or pharmacist.

Swallow the tablets with a **glass of water**.

You are advised **not to drink alcohol** with this medicine.

Doses

Depression:

Adults – 25mg three times a day increasing to 150mg-200mg a day in divided doses. In severe cases (treated in hospital) the dose may be increased up

to a maximum of 100mg three times a day.

The usual maintenance dose is between 50mg and 100mg a day in divided doses.

Elderly (over 60 years) - Initially 10mg a day increasing to 30-50mg a day.

Nightly bedwetting:

Children only, to be taken at bedtime (for no longer than 3 months and up to a maximum of 75mg a day):

Over 11 years (35-54kg) - 50-75mg a day.

8-11 years (25-35kg) - 25-50mg a day.

6-7 years (20-25kg) - 25mg a day.

Under 6 years - not recommended.

If you take more than you should

If you or the patient (or someone else) swallow a lot of tablets at the same time, or you think a child may have swallowed any, contact your nearest hospital casualty department or tell your doctor immediately. Symptoms of an overdose include fast or irregular heart beat, low blood pressure, drowsiness, fits, coma, agitation, muscle rigidity, being sick or fever.

If you forget to take the tablets

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose, take another as soon as you remember and then your next dose at the usual time.

If you stop taking the tablets

Talk to your doctor before you stop taking the tablets and follow their advice as you may experience withdrawal symptoms (see section 4).

4. Possible side effects

Like all medicines, Imipramine tablets can cause side effects, although not everybody gets them.

Stop taking the tablets and contact a doctor at once if you have the following allergic reaction, pneumonitis (fever, chills, cough, difficulty breathing, unusual weight loss, feeling

sick), a skin rash, which may be itchy, sensitivity to the sun or sun lamps, puffy, swollen face or tongue, which may be severe causing shortness of breath, shock and collapse.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

Blood: reduction in some blood cells (you may experience a sore throat, mouth ulcers and recurring infections, bleeding or bruising easily) **Endocrine system and metabolism:** disturbances in sexual function or sex drive, breast swelling in men and women, production or over-production of breast milk, changes in blood sugar levels, weight gain or loss, SIADH (syndrome of inappropriate antidiuretic hormone secretion)

Brain and central nervous system: disorientation, dizziness, tiredness or sleepiness, weakness, headache, difficulty concentrating, confusion, agitation, mood swings, aggressiveness, difficulty sleeping, delusions, seeing things that are not there, anxiety, restlessness, pins and needles, tremor, muscle spasm or lack of muscle control, speech problems, fits. Anticholinergic effects (dry mouth, constipation, blurred or double vision, sweating, hot flushes, difficulty passing water (urine), dilation of the pupil of the eye, glaucoma and blockage of the small intestine)

Heart: feeling faint when getting up (postural hypotension), high or severely low blood pressure, fast/racing heart, palpitations, irregular heart-beats, changes in ECG readings

Stomach and intestines: feeling or being sick, loss of appetite, inflammation of the mucus membranes in the mouth, tongue lesions.

Liver: impaired liver function, hepatitis, including changes in liver function (as seen in blood tests), jaundice (yellowing of the skin

and/or whites of the eyes) **Other:** hair loss, ringing in the ears, small purple red spots. An increase risk of bone fractures has been observed in patients taking this type of medicine.

Withdrawal symptoms: feeling or being sick, stomach pain, diarrhoea, difficulty sleeping, nervousness, anxiety, headache, irritability **Children:** changes in behaviour.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

5. How to store Imipramine

Keep out of the sight and reach of children. Store below 25°C in a dry place.

Do not use Imipramine tablets after the expiry date stated on the label/carton/bottle. The expiry date refers to the last day of that month.

Return any unused medicines to your pharmacist for safe disposal.

6. Further information

The active substance is imipramine hydrochloride.

Each film coated tablet contains:
Imipramine Hydrochloride USP 25mg
Excipients q.s.

The other ingredients are carnauba wax, colloidal silica, gelatin, lactose, magnesium stearate, maize starch, polyvidone, stearic acid, sodium hydroxide.

What Imipramine tablets look like and contents of the pack

Imipramine tablets are circular coated tablets. Packed in the Blister Packs.

Pack Size: 7, 14, 28, 30, 50, 90, 100 and 500 tablets.



7. Manufactured In India By:

TAJ PHARMACEUTICALS LTD.

Mumbai, India

Unit No. 214.Old Bake House,
Maharashtra chambers of Commerce Lane,
Fort, Mumbai - 400001
at:Gujarat, INDIA.

Customer Service and Product Inquiries:

1-800-TRY-FIRST (1-800-222-434 & 1-800-
222-825)

Monday through Saturday 9:00 a.m. to 7:00 p.m.

EST

E-mail: tajgroup@tajpharma.com