
PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION



Toll Free No.: 1800 222 434
& 1800 222 825

Keeping medicines affordable | At Taj Pharma
HELPING DELIVER LIFE-CHANGING THERAPIES
All trademarks mentioned enjoy legal protection.
www.tajpharma.com / www.tajoncology.com
E-mail: info@tajpharma.com / tajgroup@tajpharma.com

Submission Control No: TAJ/119



© 2019 Taj Pharmaceuticals Ltd.
Trademarks are owned by or licensed to the Taj Pharmaceuticals Ltd.

Date of Revision:
May 30th, 2019





TAJ PHARMA



250mg, 500mg, 750mg and 1000mg
FILM-COATED TABLETS

Levetiracetam TAJ PHARMA 250 mg film-coated tablets

Levetiracetam TAJ PHARMA 500 mg film-coated tablets

Levetiracetam TAJ PHARMA 750 mg film-coated tablets

Levetiracetam TAJ PHARMA 1,000 mg film-coated tablets

Read all of this leaflet carefully before you or your child start taking this medicine because it contains important information for you.

Keep this leaflet. You may need to read it again.

If you have any further questions, ask your doctor or pharmacist.

This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.

If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Levetiracetam TAJ PHARMA is and what it is used for

2. What you need to know before you take Levetiracetam TAJ PHARMA

3. How to take Levetiracetam TAJ PHARMA

4. Possible side effects

5. How to store Levetiracetam TAJ PHARMA

6. Contents of the pack and other information

1. What Levetiracetam TAJ PHARMA is and what it is used for

Levetiracetam is an antiepileptic medicine (a medicine used to treat seizures in epilepsy).

Levetiracetam TAJ PHARMA is used:

on its own in adults and adolescents from 16 years of age with newly diagnosed epilepsy, to treat a certain form of epilepsy. Epilepsy is a condition where the patients have repeated fits (seizures). Levetiracetam is used for the epilepsy form in which the fits initially affect only one side of the brain, but could thereafter extend to larger areas on both sides of the brain (partial onset seizure with or without secondary generalisation). Levetiracetam has been given to you by your doctor to reduce the number of fits.

as an add-on to other antiepileptic medicines to treat: partial onset seizures with or without generalisation in adults, adolescents, children and infants from one month of age;

myoclonic seizures (short, shock-like jerks of a muscle or group of muscles) in adults and adolescents from 12 years of age with juvenile myoclonic epilepsy;

primary generalised tonic-clonic seizures (major fits, including loss of consciousness) in adults and adolescents from 12 years of age with idiopathic generalised epilepsy (the type of epilepsy that is thought to have a genetic cause).

2. What you need to know before you take Levetiracetam TAJ PHARMA

Do not take Levetiracetam TAJ PHARMA if you are allergic to levetiracetam, pyrrolidone derivatives or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor before taking Levetiracetam TAJ PHARMA

If you suffer from kidney problems, follow your doctor's instructions. He/she may decide if your dose should be adjusted.

If you notice any slow down in the growth or unexpected puberty development of your child, please contact your doctor.

A small number of people being treated with anti-epileptics such as Levetiracetam TAJ PHARMA have had thoughts of harming or killing themselves. If you

have any symptoms of depression and/or suicidal ideation, please contact your doctor.

Children and adolescents

Levetiracetam TAJ PHARMA is not indicated in children and adolescents below 16 years on its own (monotherapy).

Other medicines and Levetiracetam TAJ PHARMA

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Do not take macrogol (a drug used as laxative) for one hour before and one hour after taking levetiracetam as this may result in a loss of its effect.

Pregnancy and breast-feeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Levetiracetam TAJ PHARMA should not be used during pregnancy unless clearly necessary. A risk of birth defects for your unborn child cannot be completely excluded. Levetiracetam has shown unwanted reproductive effects in animal studies at dose levels higher than you would need to control your seizures.

Breast-feeding is not recommended during treatment. Driving and using machines

Levetiracetam TAJ PHARMA may impair your ability to drive or operate any tools or machinery, as it may make you feel sleepy. This is more likely at the beginning of treatment or after an increase in the dose. You should not drive or use machines until it is established that your ability to perform such activities is not affected.

Levetiracetam TAJ PHARMA 750 mg tablets contain Sunset Yellow (E110) Sunset Yellow (E110) colouring agent may cause allergic reactions.

3. How to take Levetiracetam TAJ PHARMA

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Take the number of tablets following your doctor's instructions.

Levetiracetam TAJ PHARMA must be taken twice a day, once in the morning and once in the evening, at about the same time each day.

Monotherapy

Dose in adults and adolescents (from 16 years of age):

General dose: between 1,000 mg and 3,000 mg each day.

When you will first start taking Levetiracetam TAJ PHARMA, your doctor will prescribe you a lower dose (500 mg each day) during 2 weeks before giving you the lowest general dose of 1,000 mg.

Add-on therapy

Dose in adults and adolescents (12 to 17 years) weighing 50 kg or more:

General dose: between 1,000 mg and 3,000 mg each day.

Example: if your daily dose is 1,000 mg, you might take 2 tablets of 250 mg in the morning and 2 tablets of 250 mg in the evening.

Dose in infants (1 month to 23 months), children (2 to 11 years) and adolescents (12 to 17 years) weighing less than 50 kg:

Your doctor will prescribe the most appropriate pharmaceutical form of levetiracetam according to the age, weight and dose.

An oral solution is a formulation more appropriate to infants and children under the age of 6 years and to children and adolescents (from 6 to 17 years) weighing less than 50 kg and when tablets do not allow accurate dosage.

Method of administration

Swallow Levetiracetam TAJ PHARMA tablets with a sufficient quantity of liquid (e.g. a glass of water). You may take Levetiracetam TAJ PHARMA with or without food.

Duration of treatment

Levetiracetam TAJ PHARMA is used as a chronic treatment. You should continue Levetiracetam TAJ PHARMA treatment for as long as your doctor has told you.

Do not stop your treatment without your doctor's advice as this could increase your seizures.

If you take more Levetiracetam TAJ PHARMA than you should

The possible side effects of an overdose of levetiracetam are sleepiness, agitation, aggression, decrease of alertness, inhibition of breathing and coma.

Contact your doctor if you took more tablets than you should. Your doctor will establish the best possible treatment of overdose.

If you forget to take Levetiracetam TAJ PHARMA

Contact your doctor if you have missed one or more doses.

Do not take a double dose to make up for a forgotten tablet.

If you stop taking Levetiracetam TAJ PHARMA

If stopping treatment, Levetiracetam TAJ PHARMA should be discontinued gradually to avoid an increase of seizures.

Should your doctor decide to stop your Levetiracetam TAJ PHARMA treatment, he/she will instruct you about the gradual withdrawal of Levetiracetam TAJ PHARMA.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most frequently reported side effects are nasopharyngitis, somnolence (sleepiness), headache, fatigue and dizziness. At the beginning of the treatment or at dose increase side effects like sleepiness, tiredness and dizziness may be more common. These effects should however decrease over time.

Very common: may affect more than 1 in 10 people

nasopharyngitis;
somnolence (sleepiness), headache.

Common: may affect up to 1 in 10 people

anorexia (loss of appetite);
depression, hostility or aggression, anxiety, insomnia, nervousness or irritability;
convulsion, balance disorder (equilibrium disorder), dizziness (sensation of unsteadiness), lethargy (lack of energy and enthusiasm), tremor (involuntary trembling);
vertigo (sensation of rotation);
cough;
abdominal pain, diarrhoea, dyspepsia (indigestion), vomiting, nausea;
rash;
asthenia/fatigue (tiredness).
Uncommon: may affect up to 1 in 100 people

decreased number of blood platelets, decreased number of white blood cells;
weight decrease, weight increase;
suicide attempt and suicidal ideation, mental disorder, abnormal behaviour, hallucination, anger, confusion, panic attack, emotional instability/mood swings, agitation;
amnesia (loss of memory), memory impairment (forgetfulness), abnormal coordination/ataxia (impaired coordinated movements), paraesthesia (tingling), disturbance in attention (loss of concentration);
diplopia (double vision), vision blurred;
elevated/abnormal values in a liver function test;
hair loss, eczema, pruritus;
muscle weakness, myalgia (muscle pain);
injury.
Rare: may affect up to 1 in 1,000 people

infection;
decreased number of all blood cell types;
severe allergic reactions (DRESS, anaphylactic reaction [severe and important allergic reaction], Quincke's oedema [swelling of the face, lips, tongue and throat]);
decreased blood sodium concentration;
suicide, personality disorders (behavioural problems), thinking abnormal (slow thinking, unable to concentrate);
uncontrollable muscle spasms affecting the head, torso and limbs, difficulty in controlling movements, hyperkinesia (hyperactivity);
pancreatitis;
liver failure, hepatitis;

skin rash, which may form blisters and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (erythema multiforme), a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome), and a more severe form causing skin peeling in more than 30% of the body surface (toxic epidermal necrolysis).

Reporting of side effects

If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Levetiracetam TAJ PHARMA

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, label or blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Levetiracetam TAJ PHARMA contains

The active substance is called levetiracetam.

One tablet of Levetiracetam TAJ PHARMA 250 mg contains 250 mg of levetiracetam.

One tablet of Levetiracetam TAJ PHARMA 500 mg contains 500 mg of levetiracetam.

One tablet of Levetiracetam TAJ PHARMA 750 mg contains 750 mg of levetiracetam.

One tablet of Levetiracetam TAJ PHARMA 1,000 mg contains 1,000 mg of levetiracetam.

The other ingredients are:

Crospovidon, povidone, silica colloidal anhydrous, magnesium stearate, polyvinyl alcohol-part.

hydrolyzed, macrogol 4000, talc, titanium dioxide (E171), colourants*.

* The colourants are:

250 mg tablet: Indigo carmine (E132).

500 mg tablet: Iron oxide yellow (E172), indigo carmine (E132).

750 mg tablet: Indigo carmine (E132), sunset yellow (E110), iron oxide red (E172).

What Levetiracetam TAJ PHARMA looks like and contents of the pack

Levetiracetam TAJ PHARMA 250 mg film-coated tablets are oval, light blue, 13.6 x 6.4 mm marked with "L" on one side and "250" on the other side.

Levetiracetam TAJ PHARMA 500 mg tablets are oval, yellow, 17.1 x 8.1 mm marked with "L" on one side and "500" on the other side.

Levetiracetam TAJ PHARMA 750 mg tablets are oval, orange, 19.0 x 9.3 mm marked with "L" on one side and "750" on the other side.

Levetiracetam TAJ PHARMA 1,000 mg tablets are oval, white, 19.0 x 10.0 mm marked with "L" on one side and "1000" on the other side.

Perforated unit dose blisters: 60 x 1 film-coated tablets (only available for 250 mg, 500 mg and 1,000 mg tablets).

Not all pack sizes may be marketed.

Pack sizes are 28,50,100 & 500 tablets

Marketing Authorisation holder and Manufacturer:

Marketing Authorisation Holder

Manufactured in India by:

TAJ PHARMACEUTICALS LTD.

Mumbai, India

Unit No. 214.Old Bake House,
Maharashtra chambers of Commerce Lane,
Fort, Mumbai - 400001

at : Ahmedabad - Gujarat, INDIA.

Code No.: PRINTED ON BOX.

This leaflet was last revised in April 2019.



TAJ PHARMA

