## www.tajpharma.com

Norvlia® is applied daily to the nail with a bottle that has a built-in flow-through brush applicator. It dries quickly, and there is no need to remove excess product.

Norvlia is for topical use only and not for oral, ophthalmic, or intravaginal use.

For Topical Use Only (Not for use in eyes)

4 mL





Norvlia® (efinaconazole 10% topical solution) for the treatment of onychomycosis of the toenails. Norvlia® is the first topical triazole approved for the treatment of onychomycosis of the toenails.

Norvlia® (efinaconazole) topical solution, 10% For topical use

Initial U.S. Approval: 2014

## **INDICATIONS AND USAGE**

Norvlia is an azole antifungal indicated for the topical **treatment of onychomycosis** of the toenails due to Trichophyton rubrum and Trichophyton mentagrophytes.

## DOSAGE AND ADMINISTRATION

Apply Norvlia to affected toenails once daily for 48 weeks using the integrated flow-through brush applicator.

When applying Norvlia, ensure the toenail, the toenail folds, toenail bed, hyponychium, and the undersurface of the toenail plate, are completely covered.

For topical use only.

Not for oral, ophthalmic, or intravaginal use.

DOSAGE FORMS AND STRENGTHS

Solution: 10%.

CONTRAINDICATIONS: None.

ADVERSE REACTIONS

The most common adverse reactions (incidence >1%) were ingrown toenails, application site dermatitis, application site vesicles, and application site pain.

Norvlia (efinaconazole) topical solution, 10% contains 100 mg of efinaconazole in each gram of clear, colorless to pale yellow solution.



