

www.tajpharma.com

Ranolazine, sold under the trade name Pipragin by Taj Pharma, is a drug to treat angina that was first approved in 2013.

An Anti-Anginal Medication

new **Pipragin[®] ER** *Redefining Possible pain Solution*
(ranolazine) extended 500, 750 & 1000 mg release tablets

Extra all day & night long

new **Pipragin[®] ER** *Redefining Possible pain Solution*
(ranolazine) extended 500, 750 & 1000 mg release tablets

Ranolazine is an **anti-anginal medication**.
It works by improving blood flow to help the heart work more efficiently.
Ranolazine is used to **treat chronic angina (chest pain)**. Ranolazine is not for use during an acute (emergency) attack of angina.

Pipragin is used to treat adults with angina that keeps coming back, also known as chronic angina. Learn how Pipragin can help you reach your goal of angina control. Find out what to expect while taking Pipragin and how Pipragin can help.

What is Pipragin?

Pipragin is a prescription medicine used to treat adults with angina that keeps coming back (chronic angina). Pipragin may be used with other medicines that are used for heart problems and blood pressure control.

IMPORTANT SAFETY INFORMATION

Who should not take Pipragin?

Do not take Pipragin if:

Pipragin may cause serious side effects, including

changes in the electrical activity of your heart called QT prolongation. Your doctor may do tests to check the electrical activity of your heart.

Tell your doctor right away if you feel faint, lightheaded, or feel your heart beating irregularly or fast, as these may be symptoms of QT prolongation.

Also tell your doctor if you, or a family member, have a history of a heart problem called "QT prolongation" or "long QT syndrome."

kidney failure in people who already have severe kidney problems. Your doctor may need to do tests to check how your kidneys are working.

The most common side effects of Pipragin include dizziness, headache, constipation, and nausea.

Tell your doctor if you have any side effect that bothers you or does not go away.

What else do I need to know about taking Pipragin?

Tell your doctor about all medical conditions you have and about all of the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Pipragin may affect the way other medicines work and other medicines may affect how Pipragin works. Swallow Pipragin tablets whole. Do not crush, break, or chew Pipragin tablets.

If you miss a dose of Pipragin, wait to take the next dose at your regular time. Do not take more than 1 dose at a time.

Limit the amount of grapefruit products you eat or drink. Grapefruit can cause your blood levels of Pipragin to increase.

Pipragin can cause dizziness, lightheadedness, or fainting. If you have these symptoms, do not drive a car, use machinery, or do anything that needs you to be alert.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

PIPRAGIN ® THE REGISTERED TRADEMARK (®) OF TAJ PHARMACEUTICALS LTD. MUMBAI., INDIA

Redefining Possible pain Solution

A Taj Pharma™ India Product