

Rosuvastatin 5mg/10mg/20mg/40mg film-coated tablets
Taj Pharma

Package leaflet: Information for the patient

Rosuvastatin 5mg/10mg/20mg/40mg film-coated tablets

rosuvastatin

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Rosuvastatin is and what it is used for
2. What you need to know before you take Rosuvastatin
3. How to take Rosuvastatin
4. Possible side effects
5. How to store Rosuvastatin
6. Contents of the pack and other information

1. What Rosuvastatin is and what it is used for

Rosuvastatin contains the active ingredient rosuvastatin, which belongs to a group of medicines called statins.

You have been prescribed this medicine because:

- You have a high cholesterol level. This means you are at risk from a heart attack or stroke. Rosuvastatin is used in adults, adolescents and children 6 years or older to treat high cholesterol.
 - You have been advised to take a statin, because changing your diet and doing more exercise were not enough to correct your cholesterol levels. You should continue with your cholesterol-lowering diet and exercise while you are taking Rosuvastatin.

Or

- You have other factors that increase your risk of having a heart attack, stroke or related health problems.

Heart attack, stroke and other problems can be caused by a disease called atherosclerosis. Atherosclerosis is due to build up of fatty deposits in your arteries.

Why it is important to keep taking Rosuvastatin:

- Rosuvastatin is used to correct the levels of fatty substances in the blood called lipids, the most common of which is cholesterol.
- There are different types of cholesterol found in the blood - 'bad' cholesterol (LDL-C) and 'good' cholesterol (HDL-C).
- Rosuvastatin can reduce the 'bad' cholesterol and increase the 'good' cholesterol.

- It works by helping to block your body's production of 'bad' cholesterol. It also improves your body's ability to remove it from your blood.

For most people, high cholesterol does not affect the way they feel because it does not produce any symptoms. However, if it is left untreated, fatty deposits can build up in the walls of your blood vessels causing them to narrow.

- Sometimes, these narrowed blood vessels can get blocked which can cut off the blood supply to the heart or brain leading to a heart attack or a stroke. By lowering your cholesterol levels, you can reduce your risk of having a heart attack, a stroke or related health problems.
- You need to keep taking Rosuvastatin, even if it has got your cholesterol to the right level, because it prevents your cholesterol levels from creeping up again and causing build up of fatty deposits. However, you should stop if your doctor tells you to do so, or you have become pregnant.

2. What you need to know before you take Rosuvastatin

Do not take Rosuvastatin

- If you are allergic to rosuvastatin or any of the other ingredients of this medicine (listed in section 6).
- If you are pregnant or breast-feeding. If you become pregnant while taking rosuvastatin stop taking it immediately and tell your doctor. Women should avoid becoming pregnant while taking rosuvastatin by using suitable contraception.
- If you have liver disease.
- If you have severe kidney problems.

- If you have repeated or unexplained muscle weakness, aches or pains.

- If you take a drug called ciclosporin (used, for example, after organ transplants) If any of the above applies to you (or if you are in doubt), please see your doctor.

In addition, do not take Rosuvastatin 40mg (the highest dose):

- If you have moderate kidney problems (if in doubt, please ask your doctor).
- If your thyroid gland is not working properly.
- If you have had any repeated or unexplained muscle aches or pains, a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines.
- If you regularly drink large amounts of alcohol.
- If you are of Asian origin (Japanese, Chinese, Filipino, Vietnamese, Korean or Indian).
- If you take other medicines called fibrates to lower your cholesterol.

If any of the above applies to you (or if you are in doubt), please see your doctor.

Warnings and precautions

Talk to your doctor or pharmacist before taking Rosuvastatin. - If you have problems with your kidneys.

- If you have problems with your liver.
- If you have a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines.
- If you regularly drink large amounts of alcohol. - If your thyroid gland is not working properly.

- If you take other medicines called fibrates to lower your cholesterol. Please read this leaflet carefully, even if you have taken other medicines for high cholesterol before.
- If you take medicines used to fight the HIV infection e.g. lopinavir/ritonavir and/or atazanavir, please see 'Other medicines and Rosuvastatin'.
- If you are taking or have taken in the last 7 days a medicine called fusidic acid (a medicine for bacterial infection), orally or by injection. The combination of fusidic acid and Rosuvastatin can lead to serious muscle problems (rhabdomyolysis). Please see 'Other medicines and Rosuvastatin'
- If you have severe respiratory failure.
- If you are of Asian origin - that is Japanese, Chinese, Filipino, Vietnamese, Korean or Indian. Your doctor needs to choose the right start dose of Rosuvastatin to suit you.

During treatment

Tell your doctor or pharmacist immediately if you have repeated or unexplained muscle aches or pains especially if you feel unwell or have a fever. Also tell your doctor or pharmacist if you have a muscle weakness that is constant. Additional test and medicines may be needed to diagnose and treat this.

Talk to your doctor or pharmacist if you develop a persistent cough, shortness of breath or breathing difficulties. This medicine can cause scarring of the lungs and you may need to be checked by your doctor.

Children and adolescents

- If the patient is under 6 years old: Rosuvastatin should not be

given to children younger than 6 years.

- If the patient is below 18 years of age: The Rosuvastatin 40mg tablet is not suitable for use in children and adolescents below 18 years of age.

Elderly

- If you are over 70 (as your doctor needs to choose the right start dose of Rosuvastatin to suit you)

If any of the above applies to you (or if you are not sure):

Do not take Rosuvastatin 40mg (the highest dose) and check with your doctor or pharmacist before you actually start taking any dose of Rosuvastatin.

In a small number of people, statins can affect the liver. This is identified by a simple test, which looks for increased levels of liver enzymes in the blood. For this reason, your doctor will usually carry out this blood test (liver function test) before and during treatment with Rosuvastatin.

While you are on this medicine your doctor will monitor you closely if you have diabetes or are at risk of developing diabetes. You are likely to be at risk of developing diabetes if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure.

Other medicines and Rosuvastatin

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including the following:

- ciclosporin (used for example, after organ transplants)

- warfarin or clopidogrel (or any other drug used for thinning the blood)
- fibrates (such as gemfibrozil, fenofibrate) or any other medicine used to lower cholesterol (such as ezetimibe)
- indigestion remedies (used to neutralise acid in your stomach) - erythromycin (an antibiotic)
- fusidic acid (an antibiotic - please see below and 'Warnings and precautions') - an oral contraceptive (the pill)
- regorafenib (used to treat cancer) - hormone replacement therapy
- any of the following drugs used to treat viral infections, including HIV or hepatitis C infection, alone or in combination (please see Warnings and precautions): ritonavir, lopinavir, atazanavir, simeprevir, ombitasvir, paritaprevir, dasabuvir, velpatasvir, grazoprevir, elbasvir, glecaprevir, pibrentasvir.

The effects of these medicines could be changed by Rosuvastatin or they could change the effects of Rosuvastatin.

If you need to take oral fusidic acid to treat a bacterial infection you will need to temporarily stop using this medicine. Your doctor will tell you when it is safe to restart Rosuvastatin. Taking Rosuvastatin with fusidic acid may rarely lead to muscle weakness, tenderness or pain (rhabdomyolysis). See more information regarding rhabdomyolysis in section 4.

Pregnancy and breast-feeding

Do not take Rosuvastatin if you are pregnant or breast-feeding. If you become pregnant while taking Rosuvastatin stop taking it immediately and tell your doctor. Women should avoid

becoming pregnant while taking Rosuvastatin by using suitable contraception.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Most people can drive a car and operate machinery while taking Rosuvastatin - it will not affect their ability. However, some people feel dizzy during treatment with Rosuvastatin. If you feel dizzy, consult your doctor before attempting to drive or use machines.

Rosuvastatin contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Rosuvastatin also contains the colourants Sunset yellow and Allura red, which may cause allergic reactions.

For a full list of ingredients please see section 6 of this leaflet.

3. How to take Rosuvastatin

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Usual doses in adults

If you are taking Rosuvastatin for high cholesterol:

Starting dose

Your treatment with Rosuvastatin must start with the 5mg or the 10mg dose, even if you have taken a higher dose of a different statin before. The choice of your start dose will depend upon:

- Your cholesterol level.
- The level of risk you have of experiencing a heart attack or stroke.
- Whether you have a factor that may make you more sensitive to possible side effects.

Please check with your doctor or pharmacist which start dose of Rosuvastatin will best suit you.

Your doctor may decide to give you the lowest dose (5mg) if:

- You are of Asian origin (Japanese, Chinese, Filipino, Vietnamese, Korean or Indian).
- You are over 70 years of age.
- You have moderate kidney problems.
- You are at risk of muscle aches and pains (myopathy).

Increasing the dose and maximum daily dose

Your doctor may decide to increase your dose. This is so that you are taking the amount of Rosuvastatin that is right for you. If you started with a 5mg dose, your doctor may decide to double this to 10mg, then 20mg and then 40mg if necessary. If you started on 10mg, your doctor may decide to double this to 20mg and then 40mg if necessary. There will be a gap of four weeks between every dose adjustment.

The maximum daily dose of Rosuvastatin is 40mg. It is only for patients with high cholesterol levels and a high risk of heart attacks or stroke whose cholesterol levels are not lowered enough with 20mg.

If you are taking Rosuvastatin to reduce your risk of having a heart attack, stroke or related health problems:

The recommended dose is 20mg daily. However, your doctor may decide to use a lower dose if you have any of the factors mentioned above.

Use in children and adolescents aged 6-17 years

The dose range in children and adolescents aged 6 to 17 years is 5 to 20mg once daily. The usual start dose is 5mg per day and your doctor may gradually increase this to find the right amount of Rosuvastatin for you. The maximum daily dose of Rosuvastatin is 10mg or 20mg for children aged 6 to 17 years depending on the underlying condition being treated. Take your dose once a day. Rosuvastatin 40mg tablet should not be used by children.

Taking your tablets

Swallow each tablet whole with a drink of water.

Take Rosuvastatin once daily. You can take it at any time of the day with or without food.

Try to take your tablet at the same time every day to help you to remember it.

Regular cholesterol checks

It is important to go back to your doctor for regular cholesterol checks, to make sure your cholesterol has reached and is staying at the correct level.

Your doctor may decide to increase your dose so that you are taking the amount of Rosuvastatin that is right for you.

If you take more Rosuvastatin than you should Contact your doctor or nearest hospital for advice.

If you go into hospital or receive treatment for another condition, tell the medical staff that you're taking Rosuvastatin.

If you forget to take Rosuvastatin

Don't worry; just take your next scheduled dose at the correct time. Do not take a double dose to make up for a forgotten dose.

If you stop taking Rosuvastatin

Talk to your doctor if you want to stop taking Rosuvastatin. Your cholesterol levels might increase again if you stop taking Rosuvastatin.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

It is important that you are aware of what these side effects may be. They are usually mild and disappear after a short time, however some can be serious and require medical attention.

Stop taking Rosuvastatin and seek medical help immediately if you have any of the following allergic reactions:

- Difficulty in breathing, with or without swelling of the face, lips, tongue and/or throat
- Swelling of the face, lips, tongue and/or throat, which may cause difficulty in swallowing - Severe itching of the skin (with raised lumps).

Also, stop taking Rosuvastatin and talk to your doctor immediately if you have any unusual weakness, aches or pains in your muscles which go on for longer than you might expect.

Muscle symptoms are more common

in children and adolescents than in adults. As with other statins, a very small number of people have experienced unpleasant muscle effects and rarely these have gone on to become a potentially life threatening muscle damage known as rhabdomyolysis.

You may also notice the following side effects, which could be signs of a serious problem. Seek medical help immediately if you see any of these:

- Feeling very tired or thirsty, urinating more often than usual, particularly at night – these may be signs of diabetes. This is more likely if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure. Your doctor will monitor you while you are taking this medicine.
- A severe stomach pain that spreads to the back (may be signs of an inflamed pancreas)
- Bruising or bleeding more often or more easily than usual, which may be signs of a low number of platelets in the blood (thrombocytopenia)
- Yellowing of the skin and eyes (jaundice), with fever, stomach pain, pale stools or dark urine (may be signs of problems with your liver)
- Feelings of numbness, tingling or burning in the arms or legs (may be signs of damage to the nerves)
- Serious skin reactions causing blistering and peeling, especially around the mouth, eyes and genitals (Stevens-Johnson syndrome)
- Breathing problems, including persistent cough and/or shortness of breath or fever (may

- be signs of scarring on the lungs, known as interstitial lung disease)
- Lupus-like disease syndrome (including rash, joint disorders and effects on blood cells)
 - Muscle rupture (muscle damage due to rupture of muscle fibres; symptoms may include pain, swelling, bruising, tenderness and loss of function).

Other possible side effects

Common (may affect up to 1 in 10 people): -
 Headache

- Stomachpain
- Constipation
- Feeling sick
- Muscle pain
- Feeling weak
- Dizziness

- An increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking your Rosuvastatin tablets (only Rosuvastatin 40mg)

Uncommon (may affect up to 1 in 100 people):

- Rash, itching, itchy rash with raised lumps (hives)
- An increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking your Rosuvastatin tablets (only Rosuvastatin 5mg, 10mg and 20mg).

Rare (may affect up to 1 in 1,000 people):

- Increase in liver enzymes, as seen in blood tests - .

Very rare (may affect up to 1 in 10,000 people): -
 Traces of blood in your urine

- Joint pain
- Memory loss
- Breast enlargement in men (gynaecomastia)

Not known (frequency cannot be estimated from the available data): - Diarrhoea (loose stools)

- Cough
- Shortness of breath - Swelling (oedema)
- Sleep disturbances, including insomnia and nightmares - Sexual difficulties
- Depression
- Tendon disorders, sometimes complicated by breaking or tearing

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

5. How to store Rosuvastatin

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the box/blisters/label after EXP. The expiry date refers to the last day of that month.
- Store below 25°C. Store in the original package in order to protect from moisture.

- Rosuvastatin supplied in bottle packs should be used for no longer than 3 months after the bottle has first been opened.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Industrial Estate, At & Post-Moraiya,
Tal-Sanand, Dist- Ahmedabad Gujarat (India)

6. Contents of the pack and other information

What Rosuvastatin contains

The active substance is rosuvastatin.

Each tablet contains either 5mg rosuvastatin (as rosuvastatin calcium) The other ingredients are:

Tablet core:

Microcrystalline cellulose, lactose monohydrate, colloidal anhydrous silica, crospovidone, magnesium oxide, magnesium stearate, iron oxide red

Film-coating:

Lactose monohydrate, hypromellose, titanium dioxide, triacetin, Quinoline yellow

What Rosuvastatin looks like and contents of the pack

Rosuvastatin 5mg: a pink, round.

Rosuvastatin is available in blister packs of 14, 15, 28, 30, 56, 60, 84, 90 and 98 tablets or in plastic bottles containing 28, 30, 56, 60, 84 and 90 tablets.

7. Manufactured By:

Taj Pharmaceuticals Ltd.

at: Plot. No. 220, Mahagujarat