

www.tajpharma.com

Vasceos® should be taken with food  
In clinical studies, Vasceos was taken with a light meal or snack  
if more than 2 to 3 hours had passed since the evening meal.  
Vasceos® (prednisone) delayed-release tablets should  
be taken once a day

**Vasceos**®  
(Prednisone Delayed-Release Tablets)  
1 MG , 2 MG & 5 MG Rheumatoid Arthritis and  
Multiple Additional Indications

Vasceos® is designed to help reduce the pain and inflammation associated with key rheumatologic conditions, including rheumatoid arthritis and polymyalgia rheumatica. Treatment for: Rheumatoid Arthritis, Polymyalgia Rheumatica, Psoriatic Arthritis, Ankylosing Spondylitis, Asthma, Chronic Obstructive Pulmonary Disease  
Vasceos® (prednisone) is a delayed release corticosteroid indicated as an anti-inflammatory or immunosuppressive agent to treat a broad range of diseases including rheumatoid arthritis, polymyalgia rheumatica, psoriatic arthritis, ankylosing spondylitis, asthma and chronic obstructive pulmonary disease (COPD).

**Vasceos**®  
(Prednisone Delayed-Release Tablets)  
1 MG , 2 MG & 5 MG Rheumatoid Arthritis and  
Multiple Additional Indications

## FDA Approval of Vasceos® (prednisone) Delayed-Release Tablets for Rheumatoid Arthritis and Multiple Additional Indications.

Drug Administration (FDA) has approved Vasceos® (prednisone) delayed-release tablets (1 mg, 2 mg and 5 mg) to treat a broad range of diseases including rheumatoid arthritis (RA), polymyalgia rheumatica (PMR), psoriatic arthritis (PsA), ankylosing spondylitis (AS), asthma and chronic obstructive pulmonary disease (COPD)

Vasceos® (prednisone) delayed-release tablets can be conveniently timed to work when your disease is most active

Vasceos releases prednisone about 4 hours after you take the tablet

You may already be familiar with prednisone. It is recognized as a medication that reduces the pain and inflammation of conditions like rheumatoid arthritis (RA) and polymyalgia rheumatica (PMR).

Vasceos works like traditional prednisone but is different because it is a delayed-release form.

This means the prednisone dose releases about 4 hours after you take Vasceos

In clinical studies, Vasceos was taken at about 10 PM

Diseases like RA and PMR are most active in the early morning hours, which can result in morning symptoms

When taken the night before (~10 PM in clinical trials) the delayed release of prednisone in Vasceos helped improve RA symptoms and reduce morning stiffness.

Vasceos is available in 5-mg, 2-mg and 1-mg tablets.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma"India Product