


www.tajpharma.com

Xlotabin® Capecitabine Tablets USP/ IP 150 & 500mg

Xlotabin (capecitabine) is indicated for the adjuvant treatment of patients with stage III (Dukes' stage C) colon cancer.

- Xlotabin is also indicated for the first-line treatment of patients with metastatic colorectal cancer.

Antineoplastic Agent



Xlotabin Efficacy that makes a difference in the **cancer treatment landscape.**

Xlotabin 150mg & 500mg
Capecitabine Tablets USP/ IP

Xlotabin (capecitabine) is a cancer medicine that interferes with the growth of cancer cells and slows their spread in the body. Xlotabin is used to treat colon cancer, and breast or colorectal cancer that has spread to other parts of the body.



WHAT IS XLOTABIN?

Xlotabin (capecitabine) is **A CANCER MEDICINE THAT INTERFERES WITH THE GROWTH OF CANCER CELLS AND SLOWS THEIR SPREAD IN THE BODY.** XLOTABIN IS USED TO TREAT COLON CANCER, AND BREAST OR COLORECTAL CANCER THAT HAS SPREAD TO OTHER PARTS OF THE BODY.

Xlotabin is often used in combination with other cancer medications and/or radiation treatments.

HOW SHOULD I TAKE XLOTABIN?

Xlotabin is usually taken twice per day. Follow the directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Xlotabin is given in a 3-week treatment cycle, and you may only need to take the medicine during the first 2 weeks of each cycle.

Your doctor will determine how long to treat you with XLOTABIN.

Xlotabin is only part of a treatment program that may also include other medications taken on different schedules.

Follow your doctor's dosing instructions very carefully.

Xlotabin should be taken with food or within 30 minutes after eating a meal.

Take Xlotabin with a full glass (8 ounces) of water.

Call your doctor if you are sick with vomiting or diarrhea, if you are unable to eat because of stomach illness, or if you are sweating more than usual. Prolonged illness can lead to dehydration or kidney failure.

You may need frequent medical tests to be sure this medicine is not causing harmful effects. Your cancer treatments may be delayed based on the results of these tests. Xlotabin can have long lasting effects on your body.

You may need frequent medical tests for a short time after you stop using this medicine.

You must remain under the care of a doctor while you are taking XLOTABIN.

Store Xlotabin at room temperature away from moisture and heat. Keep the bottle tightly closed when not in use.

Read all patient information, medication guides, and instruction sheets provided to you. **Ask your doctor or pharmacist if you have any questions.**

Food and Drug Administration (FDA) approved .Prescription Only (POM)

XLOTABIN® THE REGISTERED TRADEMARK (®) OF TAJ PHARMACEUTICALS LTD. MUMBAI., INDIA

For the treatment of patients with metastatic colorectal cancer

A Taj Pharma India Product